

AD-A034 120

AEROSPACE CORP EL SEGUNDO CALIF IVAN A GETTING LABS F/G 20/8
RELATION BETWEEN BOUNCE-AVERAGED COLLISIONAL TRANSPORT COEFFICI--ETC(U)
DEC 76 M SCHULZ

UNCLASSIFIED

TR-0077(2260-20)-15

SAMSO-TR-76-104

NL


[OF]
ADAO34120

100-1000



END

DATE
FILMED
2 - 77



ADA034120

**Relation Between Bounce-Averaged Collisional
Transport Coefficients for Geomagnetically
Trapped Electrons**

Space Sciences Laboratory
Ivan A. Getting Laboratories
The Aerospace Corporation
El Segundo, Calif. 90245

8 December 1976

Interim Report

APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED

Prepared for
SPACE AND MISSILE SYSTEMS ORGANIZATION
AIR FORCE SYSTEMS COMMAND
Los Angeles Air Force Station
P.O. Box 92960, Worldway Postal Center
Los Angeles, Calif. 90009

DDC
RECEIVED
JAN 6 1977
L

This report was submitted by The Aerospace Corporation, El Segundo, CA 90245, under Contract F04701-76-C-0077 with the Space and Missile Systems Organization, Deputy for Advanced Space Programs, P.O. Box 92960, Worldway Postal Center, Los Angeles, CA 90009. It was reviewed and approved for The Aerospace Corporation by G. A. Paulikas, Director, Space Sciences Laboratory. Lieutenant R. C. Lawson, SAMSO/YAPT, was the project engineer.

This report has been reviewed by the Information Office (OI) and is releasable to the National Technical Information Service (NTIS). At NTIS, it will be available to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication. Publication of this report does not constitute Air Force approval of the report's findings or conclusions. It is published only for the exchange and stimulation of ideas.

FOR THE COMMANDER


R. C. Lawson
1st Lt, U.S. Air Force
Technology Plans Division
Deputy for Advanced Space Programs

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

19 REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER SAMSQ-TR-76-104	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) RELATION BETWEEN BOUNCE-AVERAGED COLLISIONAL TRANSPORT COEFFICIENTS FOR GEOMAGNETICALLY TRAPPED ELECTRONS.	5. TYPE OF REPORT & PERIOD COVERED Interim rept.	6. PERFORMING ORG. REPORT NUMBER TR-0077(2260-20)-15
7. AUTHOR(s) Michael/Schulz	8. CONTRACT OR GRANT NUMBER(s) F04701-76-C-0077	9. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 12 16p.
9. PERFORMING ORGANIZATION NAME AND ADDRESS The Aerospace Corporation El Segundo, Calif. 90245	10. REPORT DATE 8 Dec 1976	11. NUMBER OF PAGES 10
11. CONTROLLING OFFICE NAME AND ADDRESS Space and Missile Systems Organization Air Force Systems Command Los Angeles, Calif. 90009	12. SECURITY CLASS. (of this report) Unclassified	13. DECLASSIFICATION/DOWNGRADING SCHEDULE
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		
15. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. T		
16. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
17. SUPPLEMENTARY NOTES		
18. KEY WORDS (Continue on reverse side if necessary and identify by block number) Fokker-Planck Equation Pitch-Angle Diffusion Collisional Transport Trapped Radiation cos squared alpha		
19. ABSTRACT (Continue on reverse side if necessary and identify by block number) The unweighted bounce average $\langle N_j \rangle$ of the atmospheric density (N_j) along a field line and the weighted bounce average $\langle (B_0/B) N_j \cos^2 \alpha \rangle$ are related (for each atmospheric constituent(j) by an integral or derivative with respect to the sine (y) of the equatorial pitch angle (α_0), where $B/B_0 (= y^{-2} \sin^2 \alpha)$ is the ratio of local to equatorial magnetic-field intensity and α is the local pitch angle. The unweighted bounce average $\langle N_j \rangle$ is relevant to energy deposition by radiation-belt particles of all species, while the weighted bounce average $\langle (B_0/B) N_j \cos^2 \alpha \rangle$ is relevant to energy deposition by alpha particles. alpha		

DD FORM 1473
(FACSIMILE)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

409 944

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

19. KEY WORDS (Continued)

cos squared alpha

20. ABSTRACT (Continued)

$\langle (B_0/B) N_{\alpha}^2 \cos^2 \alpha \rangle$ is relevant to the description of pitch-angle diffusion of radiation-belt electrons.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

PREFACE

The author is pleased to thank Dr. G. T. Davidson for a preprint introducing (7) as a new approximation for $T(y)$, and Dr. H. H. Hilton for some empirical data on $\langle N_j \rangle$.

ACCESSION for	
NTIS	NTIS Section <input checked="" type="checkbox"/>
DDC	DDC Section <input type="checkbox"/>
UNANNOUNCED	UNANNOUNCED <input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY NOTES	
DATE	FILE
A	

CONTENTS

PREFACE	1
INTRODUCTION	5
APPLICATIONS	7
REFERENCES	13

INTRODUCTION

The interaction of geomagnetically trapped electrons with the neutral atmosphere entails both a diffusion in local pitch angle (α) and a loss in electron energy (E). The corresponding transport coefficients $D_{\alpha\alpha}$ and $(dE/dt)_v$ can be constructed by summing the contributions $D_{\alpha\alpha}^{(j)}$ and $(dE/dt)_j$ from the various atmospheric constituents (j). The subscript v on dE/dt serves only to emphasize that the corresponding process is essentially frictional, as distinguished from stochastic. The partial transport coefficients $D_{\alpha\alpha}^{(j)}$ and $(dE/dt)_j$ are proportional to the local number density N_j of atmospheric constituent j but are independent [e.g., Walt, 1966] of the local pitch angle (α). Both decrease in magnitude as the energy (E) of the incident electron increases.

Radiation-belt evolution is conventionally described [e.g., Haerendel, 1968] by means of a bounce-averaged Fokker-Planck equation. Thus, it proves convenient to introduce the bounce-averaged partial transport coefficients $D_{yy}^{(j)} = \langle (B_0/B) D_{\alpha\alpha}^{(j)} \cos^2 \alpha \rangle$ and $\langle (dE/dt)_j \rangle$, where

$$y = (B_0/B)^{1/2} \sin \alpha \quad (1)$$

is the sine of the equatorial pitch angle (α_0) and B_0 is the minimum

value of B (the magnetic-field intensity) along the field line (L). The bounce average of any quantity Q is given by

$$\langle Q \rangle = (\Omega_2/2\pi) \oint (Q/v) \sec \alpha \, ds, \quad (2)$$

where v is the speed of the electron, s is the coordinate that measures arc length along B , and $2\pi/\Omega_2$ is the full bounce period defined by setting $Q = \langle Q \rangle = 1$ in (2). The purpose of the present note is to derive an analytical relationship between the contributions of a given atmospheric constituent (j) to D_{yy} and $\langle (dE/dt)_v \rangle$, i.e., to derive an analytical relationship between $\langle (B_0/B) N_j \cos^2 \alpha \rangle$ and $\langle N_j \rangle$.

DERIVATION

The essential step in the derivation is to observe that (1) implies $\cos \alpha = [1 - y^2 (B/B_0)]^{1/2}$. Thus, it follows from (2) that

$$(2\pi/\Omega_2) \langle (B_0/B) N_j \cos^2 \alpha \rangle = \oint \frac{(N_j/v)(B_0/B) \, ds}{[1 - y^2 (B/B_0)]^{-1/2}} \quad (3)$$

while

$$y(2\pi/\Omega_2) \langle N_j \rangle = \oint \frac{y (N_j/v) \, ds}{[1 - y^2 (B/B_0)]^{1/2}}. \quad (4)$$

Moreover, the right member of (4) is equal to minus the derivative of the right member of (3) with respect to y . Thus, it follows that

$$\langle (B_0/B) N_j \cos^2 \alpha \rangle = \Omega_2(y) \int_y^1 [u/\Omega_2(u)] \langle N_j \rangle \, du, \quad (5)$$

since $\cos^2 \alpha \approx 0$ for $y = 1$. This is the desired result. The practical significance of (5) is that, given either $\langle N_j \rangle$ or $\langle (B_0/B) N_j \cos^2 \alpha \rangle$ as a function of y , one can immediately obtain the other by integration or differentiation (respectively).

It can be shown by Jacobian methods [Haerendel, 1968] that the bounce-averaged phase-space density \bar{f} satisfies the Fokker-Planck equation [cf. Walt, 1966]

$$\frac{\partial \bar{f}}{\partial t} = \frac{\Omega_2}{y} \frac{\partial}{\partial y} \left[\frac{y D_{yy}}{\Omega_2} \frac{\partial \bar{f}}{\partial y} \right] - \frac{1}{y^2 v} \frac{\partial}{\partial E} \left[y^2 v \left\langle \left(\frac{dE}{dt} \right)_v \right\rangle \bar{f} \right] \quad (6)$$

in the absence of range straggling ($D_{EE} = 0$) and radial diffusion ($D_{LL} = 0$), where y is the ratio of relativistic mass (m) to rest mass (m_0). The present work has shown that the contributions of any atmospheric constituent (j) to D_{yy} and $\langle (dE/dt)_v \rangle$ in (6) are related by virtue of (5).

APPLICATIONS

The bounce period ($2\pi/\Omega_2$) in a dipolar magnetic field (B) is proportional, at fixed energy (E) and shell parameter (L), to a function called $T(y)$ which is well approximated [Davidson, 1976] by the formula

$$\begin{aligned} T(y) &\approx T(0) - [T(0) - T(1)]y^{3/4} \\ &\approx 1.380173 - 0.639693 y^{3/4}. \end{aligned} \quad (7)$$

Earlier approximations of $T(y)$ had been given by Hamlin et al. [1961] and Lenchek et al. [1961]. Although $T(y)$ can be generated to arbitrary accuracy by a numerical integration of (2) for $Q = \langle Q \rangle = 1$, the use of (7) in (5) should make the required integral for $\langle (B_0/B) N_j \cos^2 \alpha \rangle$ much easier to perform in most instances.

A particularly simple case utilizing (7) is that of a spatially uniform atmospheric density N_j . This occurs in most laboratory devices, for example. (A laboratory device that simulates the geometry of the magnetosphere is commonly called a "terrella.") In situations such as this, for which $\langle N_j \rangle = N_j$ (a constant), it follows from (5) that

$$\langle (B_0/B) N_j \cos^2 \alpha \rangle = [Z(y)/T(y)] N_j, \quad (8)$$

where

$$\begin{aligned} Z(y) &= \int_y^1 u T(u) du \\ &\approx (1/2)(1 - y^2) T(0) \\ &\quad - (4/11)(1 - y^{11/4}) [T(0) - T(1)] \end{aligned} \quad (9)$$

if $T(y)$ is approximated by (7). The present approximation yields $Z(0) \approx 0.457471$, which agrees well with the exact result [Schulz, 1974] that $Z(0) = 16/35$ (≈ 0.457143). An earlier approximation of $Z(y)$, which yielded $Z(0) \approx 0.455533$, had been derived by Schulz [1974] in a different context. It follows from (6) that the Fokker-Planck

equation for scattering by a uniform atmosphere is of the form

$$\frac{\partial \bar{f}}{\partial t} = \frac{D_{\alpha\alpha}}{y T(y)} \frac{\partial}{\partial y} \left[y Z(y) \frac{\partial \bar{f}}{\partial y} \right] - \frac{1}{y^2 v} \frac{\partial}{\partial E} \left[y^2 v \left\langle \left(\frac{dE}{dt} \right)_v \right\rangle \bar{f} \right], \quad (10)$$

where $D_{\alpha\alpha}$ and $\langle (dE/dt)_v \rangle$ are independent of y and of L .

In the real magnetosphere, of course, one does not encounter anything like a uniform atmosphere. However, various investigators have numerically implemented the bounce averages indicated in (3) and (4). The dashed and solid curves [Walt, 1966] shown in Figure 1 represent (respectively) the quantities $(2 y^2 v^3 / c^3 x^2) D_{yy}$ and $(-v/m_0 c^3) \langle (dE/dt)_v \rangle$, where x is the cosine of the equatorial pitch angle (hence $x^2 + y^2 = 1$) and c is the speed of light. These particular forms serve to factor out the energy dependence of the transport coefficients, except for some slowly varying logarithms which are evaluated at $E \sim 1$ MeV.

Except for the fact that the neutral atmosphere has several constituents (j), one should expect to find

$$\begin{aligned} & (v/m_0 c^3) \langle (dE/dt)_v \rangle \\ &= \frac{A}{y T(y)} \frac{d}{dy} \left\{ (1 - y^2) T(y) \left[\frac{2 y^2 v^3}{x^2 c^3} D_{yy} \right] \right\}, \quad (11) \end{aligned}$$

where A is a dimensionless constant that depends in detail upon the ratio between the inelastic and elastic cross sections of atmospheric molecules or atoms for incident electrons. The "data" points (filled




Figure 1. Magnitudes of $(-v/m_0c^3) \langle (dE/dt)_v \rangle$ (solid curve) and $(2\gamma^2v^3/c^3x^2) D_{yy}$ (dashed curve) for inner-zone electrons subjected solely to atmospheric collisions [Walt, 1966], together with "data" points derived from dashed curve (and nominally "predicting" the solid curve) by means of (11).

circles) in Figure 1 serve as an illustration of this idea. The square-bracketed factor in (11) has been differentiated numerically at $L = 1.185$ and treated as a constant (for $x < 0.86$) at $L = 1.9$ for this purpose.

The resulting "data" points (normalized by evaluating A to force agreement at $x = 0$) follow the solid curves rather well. One would not have expected perfect agreement, since the densities (N_j) of the various atmospheric constituents (j) have different altitude profiles and contribute to D_{yy} and $\langle (dE/dt)_v \rangle$ with different weights (essentially as Z_j^2 to D_{yy} and as Z_j to $\langle (dE/dt)_v \rangle$, where Z_j is the nuclear charge). Thus, the transition from hydrogen dominance to oxygen dominance along field lines on the $L = 1.9$ drift shell may help to account for the departure of the "data" points from the solid curve at large x .

Bounce-averaged atmospheric densities $\langle N_j \rangle$ have been compiled by several investigators [e.g., Cornwall et al., 1965] in the course of studies on radiation-belt protons (for which atmospheric pitch-angle diffusion is an unimportant process). The present result, as summarized by (5), enables one to extract the weighted bounce average relevant to electron pitch-angle diffusion directly from such compilations of $\langle N_j \rangle$. For example, if one has found that $\langle N_j \rangle = \sum_n N_n^{(j)} y^{-n}$ for some range of L values, then it follows from (5) and (7) that

$$\begin{aligned} & \langle (B_0/B) N_j \cos^2 \alpha \rangle \\ &= \sum_n \frac{N_n^{(j)}}{T(y)} \left[\frac{1 - y^{2-n}}{(2-n)} - \frac{1 - y^{2.75-n}}{0.53939 (11-4n)} \right] \end{aligned} \quad (12)$$

Terms corresponding to $n = 2$ and $n = 11/4$, if present in (12), reduce to logarithmic form upon expansion. Applications such as this may enable the present discovery (which seems just mildly curious at first sight) to be useful in practice as well as in theory.

REFERENCES

- Cornwall, J. M., A. R. Sims, and R. S. White, Atmospheric density experienced by radiation-belt protons, J. Geophys. Res., **70**, 3099, 1965.
- Davidson, G. T., An improved empirical description of the bounce motion of trapped particles, J. Geophys. Res., **81**, 0000, 1976.
- Haerendel, G., Diffusion theory of trapped particles and the observed proton distribution, in Earth's Particles and Fields, edited by B. M. McCormac, p. 171, Reinhold, New York, 1968.
- Hamlin, D. A., R. Karplus, R. C. Vik, and K. M. Watson, Mirror and azimuthal drift frequencies for geomagnetically trapped particles, J. Geophys. Res., **66**, 1, 1961.
- Lenchek, A. M., S. F. Singer, and R. C. Wentworth, Geomagnetically trapped electrons from cosmic ray albedo neutrons, J. Geophys. Res., **66**, 4027, 1961.
- Schulz, M., Particle lifetimes in strong diffusion, Astrophys. Space Sci., **31**, 37, 1974.
- Walt, M., Loss rates of trapped electrons by atmospheric collisions, in Radiation Trapped in the Earth's Magnetic Field, edited by B. M. McCormac, p. 337, Reidel, Dordrecht, 1966.

THE IVAN A. GETTING LABORATORIES

The Laboratory Operations of The Aerospace Corporation is conducting experimental and theoretical investigations necessary for the evaluation and application of scientific advances to new military concepts and systems. Versatility and flexibility have been developed to a high degree by the laboratory personnel in dealing with the many problems encountered in the nation's rapidly developing space and missile systems. Expertise in the latest scientific developments is vital to the accomplishment of tasks related to these problems. The laboratories that contribute to this research are:

Aerophysics Laboratory: Launch and reentry aerodynamics, heat transfer, reentry physics, chemical kinetics, structural mechanics, flight dynamics, atmospheric pollution, and high-power gas lasers.

Chemistry and Physics Laboratory: Atmospheric reactions and atmospheric optics, chemical reactions in polluted atmospheres, chemical reactions of excited species in rocket plumes, chemical thermodynamics, plasma and laser-induced reactions, laser chemistry, propulsion chemistry, space vacuum and radiation effects on materials, lubrication and surface phenomena, photo-sensitive materials and sensors, high precision laser ranging, and the application of physics and chemistry to problems of law enforcement and biomedicine.

Electronics Research Laboratory: Electromagnetic theory, devices, and propagation phenomena, including plasma electromagnetics; quantum electronics, lasers, and electro-optics; communication sciences, applied electronics, semiconducting, superconducting, and crystal device physics, optical and acoustical imaging; atmospheric pollution; millimeter wave and far-infrared technology.

Materials Sciences Laboratory: Development of new materials; metal matrix composites and new forms of carbon; test and evaluation of graphite and ceramics in reentry; spacecraft materials and electronic components in nuclear weapons environment; application of fracture mechanics to stress corrosion and fatigue-induced fractures in structural metals.

Space Sciences Laboratory: Atmospheric and ionospheric physics, radiation from the atmosphere, density and composition of the atmosphere, aurorae and airglow; magnetospheric physics, cosmic rays, generation and propagation of plasma waves in the magnetosphere; solar physics, studies of solar magnetic fields; space astronomy, x-ray astronomy; the effects of nuclear explosions, magnetic storms, and solar activity on the earth's atmosphere, ionosphere, and magnetosphere; the effects of optical, electromagnetic, and particulate radiations in space on space systems.

THE AEROSPACE CORPORATION
El Segundo, California

• • •